

Exclusion Statistics of Composite Fermions

Piotr Sitko

Institute of Physics, Wrocław University of Technology,
Wybrzeże Wyspiańskiego 27, 50-370 Wrocław, Poland.

Abstract

The exclusion statistics parameter of composite fermions is determined as an odd number ($\alpha = 3, 5, \dots$). The statistics of composite fermion excitations at $\nu = \frac{n}{2pn+1}$ is rederived as $\alpha_{qe}^{CF} = 1 + \frac{2p}{2pn+1}$, $\alpha_{qh}^{CF} = 1 - \frac{2p}{2pn+1}$. The duality $\frac{1}{\alpha_{qe}(n, 2p)} = \alpha_{qh}(n+1, 2p)$ is found. The distribution function for $\alpha = 3$ is obtained.

Introduction

The exclusion statistics was proposed by Haldane [1] as generalization of Pauli exclusion principle. The idea of Haldane is to define the change in the available particle space when particles are added to the system (or removed):

$$\Delta d_i = - \sum_j \alpha_{ij} \Delta N_j . \quad (1)$$

The generalization of the Haldane exclusion statistics was proposed by Wu [2] and others [3, 4, 5]. The idea of generalization is to divide the available particle space for smaller cells (let us say with k states). The number of many-particle states is given by:

$$\prod_i \binom{d_{N_i} + N_i - 1}{N_i} \quad (2)$$

where $d_{N_i} = k - \alpha(N_i - 1)$, $\alpha = \alpha_{ii}$.

Johnson and Canright used the spherical geometry to find the statistics of Laughlin quasiparticles [6]. Analyzing numerical results they found the number of many-particle states of quasiholes (of the Laughlin $1/m$ state, m – odd number):

$$\binom{N_e + N_{qh}}{N_{qh}} \quad (3)$$

where N_{qh} is the number of quasiholes, $2S + 1 = m(N_e - 1) + 1 - N_{qh}$, $2S$ is the number of flux quanta piercing the sphere. According to the definition (2) $\alpha_{qh} = \frac{1}{m}$ [6]. The corresponding statistics parameter of Laughlin quasielectrons is $\alpha_{qe} = 2 - \frac{1}{m}$. Here, we perform similar analysis in order to determine the exclusion statistics of composite fermions.

Composite fermions

The exact diagonalization results for the sphere can be interpreted in terms of composite fermions [7] if the effective field $2S^* = 2S - 2p(N_e - 1)$ is introduced. The composite fermion approach predicts the same angular momentum shell for quasiparticles [7], however, the main role play composite fermions (the number of composite fermions equals the number of electrons $N_{CF} = N_e$). The Eq. (3) is:

$$\binom{2S^* + 1}{N_{qh}} \quad (4)$$

and equals

$$\begin{aligned} & \binom{N_{CF} + 2S - m(N_{CF} - 1)}{N_{CF}} \\ &= \binom{2S + 1 + (1 - m)(N_{CF} - 1)}{N_{CF}}. \end{aligned} \quad (5)$$

According to the definition (2) the statistics parameter of composite fermions $\alpha_{CF} = m$ (odd number $\alpha_{CF} = 3, 5, \dots$, for $\alpha = 1$ one has fermions).

Wu *et al.* [2] found the distribution function of α -particles. The duality between α -particles of $\alpha = 1/m$ (holes) and $\alpha = m$ was noticed by Nayak and Wilczek [3]. Our analysis reflects the duality between Laughlin quasiholes and composite fermions.

The distribution function is given by the set of equations [2]:

$$n_i = \frac{1}{w + \alpha} \quad (6)$$

$$w^\alpha(1 + w)^{1-\alpha} = \xi \quad (7)$$

$$\xi = e^{\frac{\epsilon_i - \mu}{kT}} . \quad (8)$$

For example, for $\alpha = 3$ the solution is

$$w = \frac{1}{s_+ + s_- - \frac{2}{3}} \quad (9)$$

where

$$s_{\pm} = \left[\frac{1}{2\xi} + \frac{1}{27} \pm \sqrt{\frac{1}{\xi} \left(\frac{1}{\xi} + \frac{1}{27} \right)} \right]^{\frac{1}{3}} . \quad (10)$$

It is interesting to consider also the composite fermion excitations. Let us consider the filling $\nu = \frac{n}{2pn+1}$, then one gets n filled shells (in the field $2S^* = 2S - 2p(N_e - 1)$ the degeneration of the n -th effective shell is $2S^* + 2n - 1$). When one creates N_{qe} quasielectrons (of the state $\nu = \frac{n}{2pn+1}$) in the $(n+1)$ -th level, the number of many-particle states is

$$\binom{2S_{qe}^* + 2n + 1}{N_{qe}} \quad (11)$$

and

$$2S_{qe}^* = -\frac{2p}{1 + 2pn} N_{qe} + \frac{2S - 2pn^2 + 2p}{1 + 2pn} . \quad (12)$$

If quasiholes are present in the n -th level then

$$\binom{2S_{qh}^* + 2n - 1}{N_{qh}} \quad (13)$$

and

$$2S_{qh}^* = \frac{2p}{1 + 2pn} N_{qh} + \frac{2S - 2pn^2 + 2p}{1 + 2pn} . \quad (14)$$

Hence, according to the definition (2)

$$\alpha_{qe} = 1 + \frac{2p}{1 + 2pn} , \quad (15)$$

$$\alpha_{qh} = 1 - \frac{2p}{1 + 2pn} \quad (16)$$

as was first obtained in Ref. [8]. For $n = 1$ one gets the Laughlin states $\frac{1}{m} = \frac{1}{2p+1}$. One can notice the duality between quasielectrons of the state $\frac{n}{2pn+1}$ and quasiholes of the state $\frac{n+1}{2p(n+1)+1}$:

$$\frac{1}{\alpha_{qe}(n, 2p)} = \alpha_{qh}(n+1, 2p) . \quad (17)$$

For example, consider quasielectrons at $1/3$ and quasiholes at $2/5$.

Conclusions

It is found that composite fermions can be described within the generalized exclusion statistics as particles with statistics parameter α being an odd number. Also, the exclusion statistics of composite fermion excitations is rederived as $\alpha_{qe} = 1 + \frac{2p}{1+2pn}$, $\alpha_{qh} = 1 - \frac{2p}{1+2pn}$. The duality $\frac{1}{\alpha_{qe}(n,2p)} = \alpha_{qh}(n+1, 2p)$ is found.

This work was supported by KBN grant No. 2 P03B 111 18.

References

References

- [1] F. D. M. Haldane, Phys. Rev. Lett. **67** (1991) 937
- [2] Y-S. Wu, Phys. Rev. Lett. **73** (1994) 922
- [3] C. Nayak, F. Wilczek, Phys. Rev. Lett. **73** (1994) 2740
- [4] A. K. Rajagopal, Phys. Rev. Lett. **74** (1995) 1048
- [5] A. Dasnières de Veigy, S. Ouvry, Phys. Rev. Lett. **72** (1994) 600
- [6] M. D. Johnson, G. S. Canright, Phys. Rev. B **49** (1994) 2947
- [7] X. M. Chen, J. J. Quinn, Solid State Commun. **92** (1994) 865
- [8] S. B. Isakov, G. S. Canright, M. D. Johnson, Phys. Rev. B **55** (1997) 6727